



POSTER PRESENTATION

Open Access

# NDRG2 plays a pivotal role in ATLL leukaemogenesis by regulating the PTEN-mediated PI3K/AKT signalling pathway

Shingo Nakahata<sup>1</sup>, Tomonaga Ichikawa<sup>1</sup>, Yusuke Saito<sup>1</sup>, Yasuhito Arai<sup>2</sup>, Tomohiko Taki<sup>3</sup>, Masafumi Taniwaki<sup>4</sup>, Kazuhiro Morishita<sup>1\*</sup>

From 16th International Conference on Human Retroviruses: HTLV and Related Viruses  
Montreal, Canada. 26-30 June 2013

Adult T-cell leukemia/lymphoma (ATLL) is a rare and aggressive T-cell leukemia/lymphoma that is etiologically linked to infection by human T-cell lymphotropic virus type 1 (HTLV-1). To find candidate causative genes for ATLL development, we used an integrative genomic analysis of ATLL and identified N-myc downstream-regulated gene 2 (NDRG2) as a candidate tumour suppressor gene on chromosome 14q11. Here we demonstrate that NDRG2 is a novel PTEN-interacting protein and recruits protein phosphatase 2A to facilitate the dephosphorylation of PTEN at the Ser380, Thr382, and Thr383 (STT) cluster within the C terminus. Although the PI3K/AKT signalling pathway is frequently activated in ATLL cells, neither the genetic alterations of the PTEN and PIK3CA genes nor the transcriptional down-regulation of PTEN expression were detected in ATLL. We found that activation of the PI3K/AKT signalling pathway in ATLL is mediated through hyperphosphorylation of PTEN-STT, a well-known mechanism of PTEN inactivation, by down-regulation of NDRG2 expression. Moreover, we found that NDRG2 deficiency in mice results in constitutive PI3K/AKT activation in various organs and high incidence of T-cell lymphoma and other types of cancer. Since down-regulation of NDRG2 expression via promoter methylation has been reported in various types of cancer, activation of the PI3K/AKT pathway by the enhanced PTEN-STT phosphorylation through the inactivation of NDRG2 is crucial for the development of a wide range of cancer cells including ATLL, which harbor no activating genetic alterations in the PI3K/AKT pathway.

## Authors' details

<sup>1</sup>Division of Tumor and Cellular Biochemistry, Department of Medical Sciences, University of Miyazaki, Kiyotake, Miyazaki, Japan. <sup>2</sup>Division of Cancer Genomics, National Cancer Center Research Institute, Chuo-ku, Tokyo, Japan. <sup>3</sup>Department of Molecular Diagnostics and Therapeutics, Kyoto Prefectural University of Medicine, Kamigyo-ku, Kyoto, Japan. <sup>4</sup>Department of Molecular Hematology and Oncology, Kyoto Prefectural University of Medicine, Kamigyo-ku, Kyoto, Japan.

Published: 7 January 2014

doi:10.1186/1742-4690-11-S1-P87

**Cite this article as:** Nakahata *et al.*: NDRG2 plays a pivotal role in ATLL leukaemogenesis by regulating the PTEN-mediated PI3K/AKT signalling pathway. *Retrovirology* 2014 **11**(Suppl 1):P87.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)



\* Correspondence: [kmorishi@fc.miyazaki-u.ac.jp](mailto:kmorishi@fc.miyazaki-u.ac.jp)

<sup>1</sup>Division of Tumor and Cellular Biochemistry, Department of Medical Sciences, University of Miyazaki, Kiyotake, Miyazaki, Japan  
Full list of author information is available at the end of the article