

Poster presentation

Open Access

P04-44. Generation of antibody responses to HIV-1 membrane proximal external region (MPER) antigen

TM Holl^{*1}, M Kuraoka¹, D Liao¹, L Verkoczy², MA Moody², M Alam², H Liao², BF Haynes² and GH Kelsoe¹

Address: ¹Dept. of Immunology, Duke University, Durham, NC, USA and ²Duke University, Human Vaccine Institute, Durham, NC, USA

* Corresponding author

from AIDS Vaccine 2009
Paris, France. 19–22 October 2009

Published: 22 October 2009

Retrovirology 2009, **6**(Suppl 3):P72 doi:10.1186/1742-4690-6-S3-P72

This abstract is available from: <http://www.retrovirology.com/content/6/S3/P72>

© 2009 Holl et al; licensee BioMed Central Ltd.

Background

Immune responses to HIV-1 rarely generate broadly cross-reactive, neutralizing antibodies (NAb). Some NABs are polyreactive and bind self-antigens, suggesting that effective humoral responses to some HIV-1 antigens are constrained by tolerization of NAB B cells.

Methods

The human HIV-1 NAb 2F5 was used to identify human and mouse self-antigens that cross-react with the gp41 MPER. Mouse B cells were generated absent the tolerizing environment of bone marrow (BM) in sequential cultures of non-adherent progenitor cells containing rIL-7, and later, rBAFF. RAG1-deficient mice were reconstituted with culture derived (CD) B and T cells (CD-RAG mice); 4 wks later serum immunoglobulin and autoantibody were quantified. CD-RAG and control mice were immunized with HIV-1 gp-41 MPER antigen in alum adjuvant; germinal center (GC) and antibody responses were determined 2 weeks after primary or boost immunizations.

Results

The 2F5 NAb binds to nuclear antigens in both uninfected human and mouse cells. BM and CD transitional B cells are enriched for reactivity to self and to an MPER antigen. Sera of CD-RAG and control mice have similar IgM/IgG serum levels but autoantibody titers are greatly elevated in CD-RAG animals. MPER antigen was weakly immunogenic in C57BL/6 mice, but MPER immunization of CD-

RAG mice elicited strong (4-fold higher) GC responses and ~30-fold more MPER IgG antibody.

Conclusion

The 2F5 NAb binds to phylogenetically conserved nuclear antigens, consistent with the hypothesis that HIV-1 evades humoral immunity by exploiting immunological tolerance. We show that transitional B cells generated *in vitro* are enriched for autoreactive cells which survive and reconstitute serum antibody *in vivo*. Immunization of autoimmune CD-RAG mice with an MPER antigen results in significantly higher GC and antibody responses compared to intact controls. The CD-RAG mouse is a novel model for investigating "forbidden" humoral responses to HIV-1 antigens.