



MEETING ABSTRACT

Open Access

APH-2 and Tax expression are correlated with a HTLV-2 proviral load but not with lymphocytosis

Estelle Douceron^{1,2,3}, Zhanna Kaidarova⁴, Paola Miyazato⁵, Masao Matsuoka⁵, Edward L Murphy⁴, Renaud Mahieux^{1,2,3*}

From 15th International Conference on Human Retroviruses: HTLV and Related Viruses Leuven and Gembloux, Belgium. 5-8 June 2011

The recent discovery of HBZ, an antisense protein, encoded by HTLV-1 allowed a new way of understanding how HTLV-1 induces the development of adult T cell leukemia/lymphoma (ATLL). HBZ mRNA is expressed in all HTLV-1 patients tested regardless of their clinical status. Furthermore HBZ mRNA level is positively correlated to the HTLV-1 proviral load and it involved in infected T cell proliferation. The HTLV-2 homolog of HBZ, APH-2, also represses the viral transcription from the 5' LTR. We therefore quantified APH-2 and Tax mRNA levels as well as proviral load in a series of 51 blood samples obtained from the HTLV Outcomes Study (HOST) cohort. These samples were divided in low, intermediate and high proviral load (PVL) groups. We first show that APH-2 was expressed in most (94%) samples, while Tax was expressed mostly in the high PVL group. A positive correlation was observed between PVL and Tax and between PVL and APH-2. Although lymphocytosis is commonly observed among HTLV-2 carriers, we also demonstrate that APH2, contrary to HBZ does not promote cell proliferation *in vitro*. These results were confirmed *in vivo* since we did not observe a correlation between APH-2 level and the lymphocyte count. Our results therefore demonstrate that APH-2 is frequently expressed *in vivo* in HTLV-2 carriers. However, and contrary to HBZ, APH-2 does not promote cell proliferation.

Author details

¹Equipe Oncogénèse Rétrovirale, INSERM-U758 Virologie Humaine, Lyon, cedex 07, 69364, France. ²Ecole Normale Supérieure de Lyon, Lyon, cedex 07, 69364, France. ³IFR 128 Biosciences Lyon-Gerland, Lyon, cedex, 0769364,

* Correspondence: renaud.mahieux@ens-lyon.fr

¹Equipe Oncogénèse Rétrovirale, INSERM-U758 Virologie Humaine, Lyon, cedex 07, 69364, France

Full list of author information is available at the end of the article

France. ⁴University of California San Francisco, Blood Systems Research Institute, San Francisco, California, USA. ⁵Laboratory of Virus Immunology, Institute for Virus Research, Kyoto University, Kyoto, 606-8507, Japan.

Published: 6 June 2011

doi:10.1186/1742-4690-8-S1-A184

Cite this article as: Douceron *et al.*: APH-2 and Tax expression are correlated with a HTLV-2 proviral load but not with lymphocytosis. *Retrovirology* 2011 **8**(Suppl 1):A184.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

