



POSTER PRESENTATION

Open Access

# Spontaneous immunological targeting of endogenous retrovirus K is directed against the envelope protein in rhesus macaques

Enrique J Léon<sup>1,2</sup>, Lyle T Wallace<sup>3</sup>, Francesca A Norante<sup>3</sup>, Matthew B Buechler<sup>3</sup>, Laura P Newman<sup>3</sup>, Philip A Castrovinci<sup>3</sup>, Brad R Jones<sup>4</sup>, Robert J Gifford<sup>5</sup>, Jonah B Sacha<sup>1,2\*</sup>

From *Frontiers of Retrovirology* 2011  
Amsterdam, The Netherlands. 3-5 October 2011

Endogenous retroviruses (ERVs), remnants of ancient retroviral infections, have invaded the germ line of every vertebrate species including both non-human primates and humans. Most ERVs are functionally crippled by extensive deletions, mutations and hypermethylation, leading to the view that they are inert genomic fossils. However, some ERVs retain the potential for biological activity and can produce mRNA transcripts, functional viral proteins, and even non-infectious virus particles during certain developmental and pathological processes. Although ERV gene products are implicated in the etiology of disease, their impact during host infection with exogenous retroviruses such as human and simian immunodeficiency virus (HIV/SIV) remains unknown. Here, we identify the homologue of the biologically active human ERV-K, macaque ERV-K (macERV-K), which retains intact open reading frames in the macaque genome. Indeed, we isolated a fully-spliced mRNA product encoding macERV-K Env, which contains all the structural features of a canonical retroviral Envelope protein. Furthermore, we identified robust CD8+ and CD4+ T cell responses directed at macERV-K Env in one SIV-infected Indian rhesus macaque. The macERV-K Env-specific CD4+ T cell response in this animal was larger than the entire 51V-specific CD4+ T cell response and had a uniform effector memory phenotype, indicating continual low-level antigen exposure. In support of this, we also identified antibody responses targeting macERV-K in this animal. Although SIV infection did not appear to trigger ERV Env-specific T cell responses, these data suggest that Env proteins from ERVs can be active during infections with exogenous retroviruses. The interaction of ERV Env

proteins expressed in HIV-infected cells may have major implications for the pathogenicity of HIV and warrants further investigation.

#### Acknowledgements

Project supported by NIH R21 AI087474-01A1 and Gates Foundation grant OPP 1008274GCE to JBS.

#### Author details

<sup>1</sup>Vaccine and Gene Therapy Institute, Oregon Health & Science University, Beaverton, OR, USA. <sup>2</sup>Oregon National Primate Research Center, Oregon Health & Science University, Beaverton, OR, USA. <sup>3</sup>Department of Pathology and Laboratory Medicine, University of Wisconsin - Madison, WI, USA. <sup>4</sup>Department of Immunology, University of Toronto, Toronto, ON, Canada. <sup>5</sup>Aaron Diamond AIDS Research Center, Rockefeller University, New York, NY, USA.

Published: 3 October 2011

doi:10.1186/1742-4690-8-S2-P60

Cite this article as: Léon *et al.*: Spontaneous immunological targeting of endogenous retrovirus K is directed against the envelope protein in rhesus macaques. *Retrovirology* 2011 **8**(Suppl 2):P60.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)



<sup>1</sup>Vaccine and Gene Therapy Institute, Oregon Health & Science University, Beaverton, OR, USA

Full list of author information is available at the end of the article